

e-PHENO

Monitoring plant phenology in the Western Alps

Gianluca Filippa
(and the e-Pheno team)

Aosta Valley Environmental Protection Agency
ARPA VdA

Ceresole Reale

10 September 2014 - ALPARC workshop



① Why Phenology?

② e-PHENO Project

- Overview

③ The Network

- PhenoNetwork
- NDVI database
- Webcam package

④ Concluding remarks


Phenology

- The rhythm of the seasons (Morisette et al 2009, Front. Ecol. Env.)
- It responds relatively fast to weather and climate changes
- It is easy to measure
- The study of phenology has a long tradition
- Phenology is the simplest process in which to track changes in the ecology of species in response to climate change (IPCC, 2007)
- In mountain ecosystems, phenology matters even more




Phenology in mountains

Leaf emergence in Larch in Aosta Valley can vary by some 90 days in a single year




Elevation, **Continentality**, **Aspect**, Precipitation, etc.

Phenology and Climate Change




Spring onset is occurring earlier (Jeong et al 2011, Glob. Ch. Biol.)

Phenology and Climate Change



Fall onset is occurring later (Jeong et al 2011, Glob. Ch. Biol.)

Phenology and Climate Change



Growing seasons are becoming longer (Jeong et al 2011, Glob. Ch. Biol.)

Shifts in Phenology are clear!

Most recent studies clearly demonstrated that plant phenology is shifting, and such shifts are **very unlikely** due to natural variability/fluctuation only.


Hence, scientists agree on the existence of an anthropogenic component to climate change.

Phenology is an useful tool to track the impact of climate change on natural ecosystems.



Phenology and its feedbacks

Morisette et al 2009, Front. Ecol. Env.



- Weather and climate control phenology
- Phenology feedbacks on atmosphere and the climate system
- Shifts in phenology act on community structure, hence biodiversity

Keep an eye on Phenology

But from how far?

- Field observer (precise, on single individuals, on single phases, expensive)
- Remote sensing (spatially integrated, poor spatial and time resolution, noisy signal)
- Near remote sensing (spatially integrated, high resolution, unmanned, cheap, need of some infrastructure)



Keep an eye on Phenology

But from how far?

- Field observer (precise, on single individuals, on single phases, expensive)
- Remote sensing (spatially integrated, poor spatial and time resolution, noisy signal)
- Near remote sensing (spatially integrated, high resolution, unmanned, cheap, need of some infrastructure)



Keep an eye on Phenology

But from how far?

- Field observer (precise, on single individuals, on single phases, **expensive**)
- Remote sensing (spatially integrated, poor spatial and time resolution, **noisy signal**)
- Near remote sensing (spatially integrated, high resolution, unmanned, cheap, need of some infrastructure)



Keep an eye on Phenology

But from how far?

- Field observer (precise, on single individuals, on single phases, **expensive**)
- Remote sensing (spatially integrated, poor spatial and time resolution, **noisy signal**)
- **Near** remote sensing (spatially integrated, high resolution, unmanned, **cheap, need of some infrastructure**)




What do we measure?

We decided to focus on two instruments: **NDVI** and **Webcams**.

- They consist of robust, simple sensors
- They are good indicators of the seasonal development of the vegetation cover
- We have (or are currently developing) the know-how to handle these data
- They are fairly inexpensive


NDVI: Rationale

- One or two bands sensors: NIR, R, G, B, ... (Skye)
- Upward e downward sensors connected to datalogger
- Possibility to calculate different indices (eg $NDVI = \frac{NIR-R}{NIR+R}$), PRI, ...)



NDVI seasonal trajectories can be associated to plant development and, thereby **phenology**.


Webcams: Rationale



Torgnon Larch stand, 2100 m ASL

Webcams: seasonal trajectory

From the red green and blue coordinates of a given number of pixels an Index of greenness is computed



The project

- started in 2008
- cooperation project Italy-France (Interreg Alcotra)
 - PhenoAlp (2008-2011) www.phenoalp.eu
 - e-Pheno (2012-2014) www.epheno.eu
- Italy - Aosta Valley (ARPA, Parco Naturale Mont Avic, Parco Nazionale Gran Paradiso)
- France - (CREA, Parc National des Ecrins, LECA Grenoble, Parc des Bauges)
- **field observations, sensor based observations (NDVI, webcam), schools engagement**

The project

- started in 2008
- cooperation project Italy-France (Interreg Alcotra)
 - PhenoAlp (2008-2011) www.phenoalp.eu
 - e-Pheno (2012-2014) www.epheno.eu
- Italy - Aosta Valley (ARPA, Parco Naturale Mont Avic, Parco Nazionale Gran Paradiso)
- France - (CREA, Parc National des Ecrins, LECA Grenoble, Parc des Bauges)
- field observations, sensor based observations (NDVI, webcam), schools engagement

The project


- started in 2008
- cooperation project Italy-France (Interreg Alcotra)
 - PhenoAlp (2008-2011) www.phenoalp.eu
 - e-Pheno (2012-2014) www.epheno.eu
- Italy - Aosta Valley (ARPA, Parco Naturale Mont Avic, Parco Nazionale Gran Paradiso)
- France - (CREA, Parc National des Ecrins, LECA Grenoble, Parc des Bauges)
- **field observations, sensor based observations (NDVI, webcam), schools engagement**

North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)

WEBCAM AND NDVI NETWORK

This map shows the distribution of NDVI and Webcam sensors of the network.



Map Satellite

Google Imagery ©2014 TerraMetrics Terms of Use Report a map error

ePHENO

- PHENO_NET
- PHENO_SENSE
- EDU_PHENO
- PHENO_APP

NEWS

[New research was published on Environmental Research Letters](#)
May 2013

[Watch PhenoALP presentation video!](#)
Jun 2014

[NDVI database now available](#)

North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)
- most sites installed in 2012-2013
- ecosystems: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- common protocols and set-up but different cameras
- storing and processing strategy under discussion



North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)
- most sites installed in 2012-2013
- ecosystems: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- common protocols and set-up but different cameras
- storing and processing strategy under discussion



North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)
- most sites installed in 2012-2013
- **ecosystems:** grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- common protocols and set-up but different cameras
- storing and processing strategy under discussion



North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)
- most sites installed in 2012-2013
- **ecosystems**: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- common protocols and set-up but **different cameras**
- storing and processing strategy under discussion




North Western Alps phenological network

- Webcam sensors (10 sites: 5IT, 5FR)
- most sites installed in 2012-2013
- **ecosystems**: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- common protocols and set-up but **different cameras**
- storing and processing strategy **under discussion**



NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included



NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included
- similar installation protocols but different sensors (Skye and ESE-LECA-Paris)
- most sites installed in 2010-2011
- ecosystems: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- storing and processing strategy defined (e-pheno database)

NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included
- similar installation protocols but **different sensors** (Skye and ESE-LECA-Paris)
- most sites installed in 2010-2011
- ecosystems: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- storing and processing strategy defined (e-pheno database)

NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included
- similar installation protocols but **different sensors** (Skye and ESE-LECA-Paris)
- most sites installed in **2010-2011**
- ecosystems: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- storing and processing strategy defined (e-pheno database)


NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included
- similar installation protocols but **different sensors** (Skye and ESE-LECA-Paris)
- most sites installed in **2010-2011**
- **ecosystems**: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- storing and processing strategy defined (e-pheno database)

NDVI database

- NDVI database (12 sites: 5IT, 7FR) but other 6 from LECA ready to be included
- similar installation protocols but **different sensors** (Skye and ESE-LECA-Paris)
- most sites installed in **2010-2011**
- **ecosystems**: grasslands (1800-2400 m asl) and subalpine (< 2000 m asl) larch forests
- storing and processing strategy **defined** (e-pheno database)

NDVI database




NDVI sensors comparison



NDVI sensors comparison

Torgnon Larch stand



NDVI sensors comparison

Torgnon Larch stand

Snow fall event




Image Processing package

A package in the popular software for statistical computing **R**


R Core Team (2013). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
<http://www.R-project.org/>

- Open source, cost free
- Highly flexible
- Available for download soon (1 month) at <https://r-forge.r-project.org/>



ROI (Region of Interest) definition


- User can define ROIs on a reference image clicking on ROIs vertexes



PNGP: Orvieille site

Data processing


- On each ROI for each date average RGB values are computed (but gridded analysis is under development)
- Greenness index (GI) is computed as $G/(R+G+B)$
- GI data is 'cleaned' with a sequential filtering procedure (low quality images, bad light, rainy days)
- Seasonal trajectories are modelled by means of various equations (double logistic equations)
- Phenological thresholds are extracted with various methods (fixed (i.e. half peak) or based on inflection points of derivatives)
- Uncertainty is computed bootstrapping the residuals



Orvielle 2012: Different ROIs with different trajectories

Data processing

- On each ROI for each date average RGB values are computed (but gridded analysis is under development)
- Greenness index (GI) is computed as $G/(R+G+B)$
- GI data is 'cleaned' with a sequential filtering procedure (low quality images, bad light, rainy days)
- Seasonal trajectories are modelled by means of various equations (double logistic equations)
- Phenological thresholds are extracted with various methods (fixed (i.e. half peak) or based on inflection points of derivatives)
- Uncertainty is computed bootstrapping the residuals




Torgnon Larch site: 2012

Data processing

- On each ROI for each date average RGB values are computed (but gridded analysis is under development)
- Greenness index (GI) is computed as $G/(R+G+B)$
- GI data is 'cleaned' with a sequential filtering procedure (low quality images, bad light, rainy days)
- Seasonal trajectories are modelled by means of various equations (double logistic equations)
- Phenological thresholds are extracted with various methods (fixed (i.e. half peak) or based on inflection points of derivatives)
- Uncertainty is computed bootstrapping the residuals






Filtering procedure applied to Orvieille 2012 raw data


Data processing

- On each ROI for each date average RGB values are computed (but gridded analysis is under development)
- Greenness index (GI) is computed as $G/(R+G+B)$
- GI data is 'cleaned' with a sequential filtering procedure (low quality images, bad light, rainy days)
- Seasonal trajectories are modeled by means of various equations (double logistic equations)
- Phenological thresholds are extracted with various methods (fixed (i.e. half peak) or based on inflection points of derivatives)
- Uncertainty is computed bootstrapping the residuals



The double logistic function and the extracted thresholds are as in Klosterman et al. 2014

Torgnon Larch Site



The double logistic function is as in Klosterman et al. 2014, extracted thresholds are computed according to Gu et al. 2003

The package

- Developed in collaboration with PHENOCAM (<http://phenocam.sr.unh.edu/webcam/>)
- Highly flexible (5 fitting methods, 5 thresholding approaches)
- Tested on a large database (250 site-years)
- Covering a wide range of ecosystems (from arid to alpine grasslands, from deciduous tropical to coniferous boreal forests)
- Designed to be user-friendly



Costs

- NDVI measurement system
 - The sensor itself (Skye)
 - Power supply (solar panel + battery)
 - Data logger

Total: 3-3.5k Euro

- Webcam measurement system
 - The webcam itself (Reflex Nikon)
 - Power supply
 - Data logger and case

Total: 4-5k Euro

Additional costs: biweekly/monthly visit to the site

Costs

- NDVI measurement system
 - The sensor itself (Skye)
 - Power supply (solar panel + battery)
 - Data logger

Total: 3-3.5k Euro

- Webcam measurement system
 - The webcam itself (Reflex Nikon)
 - Power supply
 - Data logger and case

Total: 4-5k Euro

Additional costs: biweekly/monthly visit to the site

Costs

- NDVI measurement system
 - The sensor itself (Skye)
 - Power supply (solar panel + battery)
 - Data logger

Total: 3-3.5k Euro

- Webcam measurement system
 - The webcam itself (Reflex Nikon)
 - Power supply
 - Data logger and case

Total: 4-5k Euro

Additional costs: biweekly/monthly visit to the site

Costs

- NDVI measurement system
 - The sensor itself (Skye)
 - Power supply (solar panel + battery)
 - Data logger

Total: 3-3.5k Euro

- Webcam measurement system
 - The webcam itself (Reflex Nikon)
 - Power supply
 - Data logger and case

Total: 4-5k Euro

Additional costs: biweekly/monthly visit to the site

Summary

- The e-Pheno project allowed to build/extend a network of NDVI and WEBCAM measurements in the North Western Alps
- A database was built to host NDVI data, and it is ready to include new sites
- A package software has been designed to process Webcam images
- Data, software and know-how are freely available, in an open-source perspective

Summary

- The e-Pheno project allowed to build/extend a network of NDVI and WEBCAM measurements in the North Western Alps
- A database was built to host NDVI data, and it is ready to include new sites
- A package software has been designed to process Webcam images
- Data, software and know-how are freely available, in an open-source perspective

Summary

- The e-Pheno project allowed to build/extend a network of NDVI and WEBCAM measurements in the North Western Alps
- A database was built to host NDVI data, and it is ready to include new sites
- A package software has been designed to process Webcam images
- Data, software and know-how are freely available, in an open-source perspective

Summary

- The e-Pheno project allowed to build/extend a network of NDVI and WEBCAM measurements in the North Western Alps
- A database was built to host NDVI data, and it is ready to include new sites
- A package software has been designed to process Webcam images
- Data, software and know-how are freely available, in an open-source perspective

Summary

- The e-Pheno project allowed to build/extend a network of NDVI and WEBCAM measurements in the North Western Alps
- A database was built to host NDVI data, and it is ready to include new sites
- A package software has been designed to process Webcam images
- Data, software and know-how are freely available, in an open-source perspective

Further steps

- Improve data intercomparison (different sensors, different set-up, etc.)
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are welcome)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites

- More info

www.epheno.eu

www.arpa.vda.it/climatechange



Further steps

- **Improve data intercomparison (different sensors, different set-up, etc.)**
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are welcome)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites
- More info
www.epheno.eu
www.arpa.vda.it/climatechange

Further steps

- Improve data intercomparison (different sensors, different set-up, etc.)
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are welcome)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites
- More info
www.epheno.eu
www.arpa.vda.it/climatechange

Further steps

- Improve data intercomparison (different sensors, different set-up, etc.)
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are **welcome**)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites
- More info
www.epheno.eu
www.arpa.vda.it/climatechange

Further steps

- Improve data intercomparison (different sensors, different set-up, etc.)
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are **welcome**)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites
- More info
www.epheno.eu
www.arpa.vda.it/climatechange

Further steps

- Improve data intercomparison (different sensors, different set-up, etc.)
- Enlarge the network: incorporate existing sites, populate the database
- Enlarge the network: include new sites (e-Pheno is finishing, but the network goes on! Parks are **welcome**)
- Increase range of ecosystem types, climate, i.e. increase the representativeness of observation network
- Provide know-how and installation instructions for new partners/sites
- More info

www.epheno.eu

www.arpa.vda.it/climatechange

